

## Optional Paediatric Vaccines

(Vaccines not covered by Family Health Service, Department of Health)

Babies can visit the Family Health Service under Department of Health for basic vaccination. You can have these optional vaccines to provide a more comprehensive protection for your baby.

Vaccine/ Age	2 m	4 m	6 m	9 m	1 yr	15 m	18 m	2 yr
Haemophilus Influenzae Type b (Hib) *								
Rotavirus * (Rotarix/ RotaTeq)								
Meningococcal B (Bexsero) ∞								
Meningococcal ACWY (Menactra) *								
Hepatitis A (Havrix) §								
Japanese B Encephalitis (Imojev) §								
Influenza *	≥ 6 months; yearly during winter (Oct to Dec)							
Human Papilloma Virus (HPV/ Gardasil 9) *	9—15yr : 2 doses; >15yr : 3 doses (applicable to boys and girls)							

\* Included in National Immunization Program in USA, Canada, UK, Europe and Australia

∞ Included in National Immunization Program in USA, Canada, UK and Europe

§ Included in National Immunization Program in Japan, China and some Asian Countries

## Infectious Diseases and Vaccines in Children

Diseases	Outcome or Consequences	Vaccinations
Diphtheria	Swollen and inflamed laryngopharynx, respiratory difficulty	DTP Vaccine
Tetanus	Muscle cramps and spasms, even death	DTP Vaccine
Pertussis	Prolonged cough, pneumonia and brain damage	
Poliomyelitis	Nervous system damage, paralysis	Polio Vaccine
Haemophilus Influenzae Type B (Hib)	Meningitis, pneumonia, septicaemia	Hib Vaccine
Hepatitis B	Hepatitis and liver cancer	Hepatitis B Vaccine
Rotavirus	Fever, vomiting, diarrhoea and dehydration	Rotavirus Vaccine
Pneumococcus (Streptococcus Pneumoniae)	Bacterial pneumonia, meningitis, otitis media and septicaemia	Pneumococcal Conjugate Vaccine
Rubella	Fetal malformation, congenital blindness and heart disease	MMRV Vaccine
Measles	Meningitis and encephalitis	MMRV Vaccine
Mumps	Meningitis, infertility if infected in adolescence	
Chickenpox	Skin rash with blisters, may cause permanent scarring	
Hepatitis A	Inflammation of liver cells, liver damage	Hepatitis A Vaccine
Meningococcus	Bacterial meningitis and septicaemia, even death	Meningococcal (ACWY) Polysaccharide Vaccine & Meningococcal B Vaccine
Japanese B Encephalitis	Epilepsy, neurological damage, cognitive and motor impairment	Japanese B Encephalitis Virus Vaccine

## 胡立輝醫生 (兒科專科醫生)

本診所為嬰孩提供所需的各種防疫注射，亦會為不同成長階段的嬰孩提供定期健康檢查及智能發展評估，確保嬰孩健康成長。

電話預約查詢: 28682000 診所網頁: [www.drchriswoo.com](http://www.drchriswoo.com)

WhatsApp: 63651828

WeChat: [dr\\_chriswoo/63651828](https://www.whatsapp.com/channel/0029va80113816303760280)

診所地址: 九龍尖沙咀廣東道5號海洋中心14樓1416室

## 兒童疫苗注射計劃

若家長決定不參予政府母嬰健康接種疫苗，嬰孩可於6周至2個月到訪本診所，直接接種疫苗。

年齡	6周 - 2個月	4個月	6個月	9個月	12個月	15個月	18個月	2歲
身體檢查								
智能發展評估								
五合一疫苗								
六合一疫苗								
口服輪狀病毒疫苗*								
四痘混合疫苗								
13價肺炎鏈球菌疫苗								
甲型肝炎疫苗								
腦膜炎雙球菌疫苗 (流腦-四價 ACWY)								
腦膜炎雙球菌疫苗 (流腦B)								
日本腦炎疫苗(乙腦)								
流感疫苗	6個月以上嬰孩，每年冬季(10-12月)注射							
人類乳頭狀病毒疫苗 (HPV)	9—15歲：兩針；>15歲：三針(男女均可)							

/ 基本需要接種之疫苗

其他建議接種之疫苗

\* 輪狀病毒疫苗有兩種，分別是服兩劑(Rotarix)及三劑(RotaTeq)，效果相約。服兩劑的嬰兒最遲要於20週要服第一劑；至於服三劑的嬰兒最遲要於24週要服第一劑。

## 兒童附加疫苗 (衛生處母嬰健康院會以外)

衛生處母嬰健康院會提供基本疫苗接種予各嬰孩，以下是本診所可為嬰孩提供健康院以外的兒童附加疫苗保護，詳情請跟我們聯絡。

疫苗 / 建議 接種年齡	2 個月	4 個月	6 個月	9 個月	1歲	15 個月	18 個月	2歲
乙型流感嗜血 桿菌 (Hib) *								
輪狀病毒 * (Rotarix/ RotaTeq)								
B 型腦膜炎雙 球菌(流腦-B) ∞								
四價 (ACWY) 腦膜炎雙球菌 (流腦-四價) *								
甲型肝炎 §								
日本腦炎(乙 腦) §								
流行性感冒 *	6 個月以上嬰孩，每年冬季 (10-12 月) 注射							
人類乳頭狀病 毒(HPV/ Gardasil 9) *	9—15 歲：兩針； >15 歲：三針 (男女均可)							

\*美國、加拿大、英國、歐洲、澳洲為常規疫苗

∞美國、加拿大、英國及歐洲為常規疫苗

§日本、中國、及部分亞洲國家為常規疫苗

## 兒童常見傳染病及疫苗

傳染病	可引至後果	疫苗
白喉	咽喉發炎，呼吸困難	注射白喉、破傷風、百日咳混合疫苗
破傷風	肌肉抽筋及痙攣，嚴重會致命	
百日咳	令病者久咳、肺炎及腦受損	
小兒麻痺	神經系統受損及麻痺	注射小兒麻痺疫苗
乙型流感嗜血 桿菌	腦膜炎、肺炎、菌血症	注射乙型流感嗜血桿菌疫苗
乙型肝炎	肝炎及肝癌	注射乙型肝炎疫苗
輪狀病毒	發燒、嘔吐和腹瀉，以致脫水	服食口服輪狀病毒疫苗
肺炎鏈球菌疫苗	細菌性肺炎、菌血症、腦膜炎及中耳炎等；嚴重可致命	注射肺炎鏈球菌疫苗
德國麻疹	懷孕婦女若感染，可致畸胎，失明及先天性心臟病	注射麻疹、德國麻疹、腮腺炎及水痘疫苗 (即四痘混合疫苗)
麻疹	腦膜炎或 腦炎	
腮腺炎	腦膜炎，青春後如受感染，可能引致不育	
水痘	皮膚全身出現紅疹及水泡，水泡結痂後或會留下疤痕	
甲型肝炎	肝臟發炎，嚴重者肝臟受損	注射甲型肝炎疫苗
腦膜炎雙球菌	敗血病和細菌性腦膜炎，引致失聰及甚至致命	注射 4 價(ACWY)腦膜炎雙球菌結合疫苗(流腦)及 B 型腦膜炎雙球菌疫苗
日本腦炎 (乙腦)	腦炎引致腦部受損，引致癲癇、認知及行動障礙	注射日本腦炎疫苗(乙腦)

## Dr Chris Woo (Specialist in Paediatrics)

To prevent different infectious diseases, our clinic will provide different types of vaccinations for children. Children coming for vaccination will be also assessed by Dr Woo to ensure their good physical health and development.

Telephone booking: 28682000

Home page: [www.drchriswoo.com](http://www.drchriswoo.com)

WhatsApp: 63651828

WeChat: [dr\\_chriswoo/63651828](https://www.whatsapp.com/channel/00299a6963651828)

Address: Suite 1416, 14/F, Ocean Centre, 5 Canton Road, Tsim Sha Tsui, Kowloon

### Healthy Baby Vaccination

If baby is not visiting Family Health Service (Department of Health) for check-up and vaccination, he/she can visit our clinic directly from 6 weeks to 2 months old to start vaccination.

Age	6 wk - 2 m	4 m	6 m	9 m	12 m	15 m	18 m	2 yr
Physical Exam								
Developmental Assessment								
DTaP+IPV/ Hib								
DTaP+IPV/ Hib+HBV								
Rotavirus vaccine*								
Measles, Mumps, Rubella & Varicella vaccine								
Pneumococcal conjugate vaccine								
Hepatitis A vaccine								
Meningococcal Polysaccharide vaccine (ACWY)								
Meningococcal B vaccine								
Japanese B Encephalitis vaccine								
Influenza vaccine	≥ 6 months; Preferably in October to December each year							
Human Papilloma Virus vaccine	9—15 years: 2 doses; >15 years: 3 doses							

Basic vaccinations

Recommended optional vaccinations

\*There are two types of Rotavirus vaccines with similar efficacy; two-dose regime (Rotarix): the first dose should be taken not later than 20 weeks; three-dose regime (RotaTeq): the first dose should be taken not later than 24 weeks.